



1 Harrier Court  
 Airfield Business Park  
 Elvington  
 York YO41 4AU  
 United Kingdom  
 Tel: +44 (0) 1904 607600  
 Fax: +44 (0) 1904 607601  
 Web: www.vet-way.com

*(Levamisol 20 % also available in soluble powder...)*

## LEVAMISOL 10 % INJECTABLE 100 ml

<b>Type</b>	Anthelmintic
<b>Composition</b>	Levamisole hydrochloride: 10 g Excipient: q.s. 100 ml
<b>Intended For</b>	Cattle, sheep, pigs and goats
<b>Uses</b>	<p>For the treatment and control of parasitic bronchitis, parasitic gastro-enteritis in cattle, sheep, pigs and goats. Infections with Ascarid worms.</p> <p>When used at recommended dose, Levamisol 10% is effective against the mature and developing immature stages of Levamisole sensitive Haemonchus, Trichostrongylus, Cooperia, Nematodirus, Bunostomum and Oesophagostomum species in the gastro-intestinal tract, and Dictyocaulus spp in the lungs. Levamisole belongs to imidazothiazoles. It is very safe to use, and its availability in the blood is very quick. It can be used for pregnant or milking females. Levamisole does not induce photosensitivity and studies have proven a non specific immunostimulant effect.</p>
<b>Dosage &amp; Administration</b>	<p>Intramuscular and Subcutaneous routes</p> <p>Cattle (IM): 7.5 ml for 100 kg bodyweight</p> <p>Sheep, goats (IM route): 0.75 ml for 10 kg bodyweight</p> <p>Pigs (SC route): 0.75 ml for 10 kg bodyweight</p> <p>Another injection can be given 8 to 15 days after the first one, if necessary.</p>
<b>Warnings</b>	Very few side effects at recommended dosage. Should not be used simultaneously or up to 14 days after treatment with products containing organophosphorus compounds. To be kept away from heat and light.
<b>Withdrawal Period</b>	<p>Meat and offal: 9 days in cattle and 14 days in sheep</p> <p>Not to be used in animals producing milk for human consumption</p>

