

Photograph not available



## MAXAZOLE 2.5 % 1L

<b>Type</b>	Anthelmintic & Flukicide
<b>Composition</b>	Albendazole Oral Suspension Albendazole: 25 mg/ml
<b>Intended for</b>	Cow, sheep, goat
<b>Uses</b>	For the control of roundworms, lungworms, tapeworms, adult liver fluke, fluke and roundworm eggs. Maxazole 2.5% is a broad spectrum anthelmintic for the control of mature and developing immature forms of gastro-intestinal roundworms, lungworms, tapeworms and adult liver fluke in sheep and cattle. Maxazole 2.5% is ovicidal and will kill fluke and roundworm eggs, therefore reducing pasture contamination
<b>Administration</b>	For oral administration only using accurately calibrated oral dosing syringes. Accurate bodyweight estimation is important. 1 ml of Maxazole 2.5% contains 25 mg of Albendazole Cattle: For the control of roundworms, lungworms, tapeworms, fluke and roundworm eggs. 7.5 mg Albendazole per kg bodyweight which is equivalent to 15 ml of Maxazole 2.5 % per 50 kg of bodyweight For the additional treatment of adult liver fluke in cattle, 10 mg of albendazole per kg bodyweight, which is equivalent to 20 ml of Maxazole 2.5 % per 50 kg bodyweight. Dosage guide (see label) Sheep/goats: For the control of roundworms, lungworms, tapeworms, fluke and roundworm eggs. 5 mg of Albendazole per bodyweight which is equivalent to 2 ml of Maxazole 2.5% per 10 kg bodyweight. For the additional treatment of adult liver fluke in sheep, 7.5 mg per kg bodyweight which is equivalent to 3 ml Maxazole 2.5% per 10 kg bodyweight. Dosage guide (see label)
<b>Contra-indications, warnings</b>	Wear suitable protective clothing including rubber gloves. Wash hands after use. Do not contaminate water ways, ponds or lakes with product or empty containers. Dispose of used containers carefully. Keep out of the reach of Children. When dosing care must be taken not to damage the pharyngeal area especially in sheep. Shake container before use.
<b>Withdrawal period</b>	Cattle can be slaughtered for human consumption after 20 days from the last treatment. Milk for human consumption can be taken from cows after 72 hours from the last treatment.  Sheep/goat can be slaughtered for human consumption after 8 days from last treatment. Do not use in sheep producing milk for human consumption.  Do not dose ewes at the "fluke & worm dose" (7.5mg/kg) when rams are introduced and up to 1 month after removing rams