



1 Harrier Court
 Airfield Business Park
 Elvington
 York YO41 4AU
 United Kingdom
 Tel: +44 (0) 1904 607600
 Fax: +44 (0) 1904 607601
 Web: www.vet-way.com

OXYTETRACYCLINE SOLUBLE POWDER 25 kg

Type	Soluble powder for oral administration of oxytetracycline
Composition	Oxytetracycline hydrochloride 50g Carrier ad 1000g
Intended For	Cattle, horse, sheep, pigs, rabbits, poultry, dogs and cats
Uses	Broad spectrum antibiotic with an effective tissue diffusion which is excreted in active form by the kidneys and in the bile Cattle, horse, sheep, pigs, rabbits, poultry, dogs and cats: infections with bacteria susceptible to Oxytetracycline Poultry: Respiratory Infections: Mycoplasma, chronic respiratory disease, infectious sinusitis, infectious coryza Digestive infections: Infectious enteritis, omphalitis of hicks, Hexamitiasis Specific infections: Avian cholera, infectious synovitis, psittacosis-ornithosis Non specific infections: infections, periods of stress (vaccinations, cold snap) which cause drops in egg production slow growth and higher mortality. In other species: Susceptible bacterial infections, viral infections
Dosage & Administration	Cattle, horse, sheep, goat, pigs, rabbits, poultry, dogs and cats: 20 mg of Oxytetracycline/kg/day (10 mg/12 hours) that is 0.4 grms of powder per kg per day Dissolve the required concentration in the drinking water or in the liquidised food. Treat for 3 days minimum.
Warnings	With herbivores oral administration of oxytetracycline should be given to milk fed animals as this will minimise any upset to the intestinal flora.
Withdrawal Period	Cattle: 10 days Pigs, chickens, turkeys: 7 days Eggs: 1 day

