



1 Harrier Court  
 Airfield Business Park  
 Elvington  
 York YO41 4AU  
 United Kingdom  
 Tel: +44 (0) 1904 607600  
 Fax: +44 (0) 1904 607601  
 Web: www.vet-way.com

# VERMECTIN 1% INJECTABLE 50 ml/10 ml

<b>Type</b>	Anthelmintic & Acaricide
<b>Composition</b>	For 50ml: Ivermectin: 0.5g Excipient: q.s. 50ml
<b>Intended For</b>	Cattle, sheep and pigs
<b>Uses</b>	Vermectin is indicated for the effective treatment and control of gastrointestinal roundworms, lungworms, eye worms, ectoparasites, warbles, mange mites and sucking lice.
<b>Dosage &amp; Administration</b>	Strict subcutaneous route, using aseptic precautions, under the loose skin in front of or behind the shoulder in cattle, in the neck in sheep and behind the ear in pigs. Cattle: 0.2 mg/kg (1ml per 50 kg bodyweight) in a single injection Sheep: 0.2mg/kg (1 ml per 50 kg bodyweight) For the control of skin diseases caused by Psoroptic mange, another injection should be given 7 days after the first injection Pigs: 0.3 mg/kg (1.5 ml per 50 kg bodyweight) in a single injection
<b>Warnings</b>	Do not inject more than 10ml per injection site. This product not to be used intramuscularly or intravenously. Vermectin can be administered to cattle, sheep and pig at any stage of pregnancy or lactation provided that the milk is not intended for human consumption. Protect from direct sunlight. Keep out of reach of children, do not eat, drink or smoke while using. Wash hands after use
<b>Withdrawal Period</b>	Meat and offal: Cattle: 35 days, sheep: 42 days, pigs: 28 days Milk for human consumption: Do not use in milking animals and pregnant females 35 days before start of lactation.

