



1 Harrier Court
 Airfield Business Park
 Elvington
 York YO41 4AU
 United Kingdom
 Tel: +44 (0) 1904 607600
 Fax: +44 (0) 1904 607601
 Web: www.vet-way.com

VETPENSTREP 20/20 INJECTABLE 100 ml

Type	Suspension of penicillin G procaine and dihydrostreptomycin sulfate
Composition	Procaine Penicillin BP 20 millions U.I. Dihydrostreptomycin Sulphate BP (Vet) 20 g Excipient q.s. 100 ml
Intended For	Cattle, horses, sheep, pigs, goats, dogs and cats
Uses	Vetpenstrep injection is indicated for use in cattle, horses, pigs, sheep, dogs and cats for the treatment of infections caused by, or associated with, organisms sensitive to Penicillin and/or Streptomycin. The combination of penicillin and dihydrostreptomycin is especially useful in the treatment of mixed infections involving both gram -positive and gram -negative organisms.
Dosage & Administration	The dose should be given once daily by deep intramuscular injection for up to three consecutive days. Cattle, horses, sheep, goat : 1 ml/20 kg. Pigs (50 kg) : 2 to 5 ml. Dogs (10 kg) : 1 to 2 ml. Cats (3kg) : 0.25 to 0.6 ml.
Warnings	Contra-indicated in known cases of hypersensitivity to Penicillins . Occasionally in sucking and fattening pigs administration of this product may cause a transient pyrexia, vomiting, shivering, listlessness and incoordination. Additionally in pregnant sows and gilts a vulval discharge which could be associated with abortion, has been reported. Care should be taken not to exceed the recommended dosage. Milk for human consumption must not be taken during treatment. Shake the container before use. Store below 25°C. Protect from light. Keep out of the reach of children. Following withdrawal of the first dose, use the product within 28 days. Discard unused material.
Withdrawal Period	Meat and offal : 21 days in cattle, 31 days in sheep and 18 days in pigs Milk: 48 hours

